

Higgs Sector with Spontaneous CP Violation in $S(3)$ Standard Model

J. E. BARRADAS-GUEVARA^{1*}, O. FÉLIX-BELTRÁN², E. RODRÍGUEZ-JÁUREGUI³

¹Fac. de Cs. Físico Matemáticas, Benemérita Universidad Autónoma de Puebla, Apdo. Postal 1152, Puebla, Pue. 72000, México.

²Fac. de Cs. de la Electrónica, Benemérita Universidad Autónoma de Puebla, Apdo. Postal 542, Puebla, Pue. 72000, México.

³Departamento de Física, Universidad de Sonora, Apdo. Postal 1626, Hermosillo, Son. 83000, México.

*Email: barradas@cfm.buap.mx

Published online: August 08, 2016,

The Author(s) 2016. This article is published with open access at www.chitkara.edu.in/publications

Abstract Conditions for spontaneous Charge-Parity (CP) violation in the scalar potential sector of general $S(3)$ Higgs-doublet model (3HDM) are analyzed. An analysis of the Higgs sector of the minimal $S(3)$ -invariant extension of the Standard Model including CP violation arising from the spontaneous breaking of the electroweak symmetry is presented. This extended Higgs sector with three $SU(2)$ doublets Higgs fields with complex vev's provides an interesting scenario to analyze the Higgs masses spectrum, trilinear self-couplings and CP violation. We present how the spontaneous electroweak symmetry breaking, coming from three $S(3)$ Higgs fields, gives an interesting scenario with nine physical Higgs and three Goldstone bosons, when spontaneous CP violation arises from the Higgs field $S(3)$ singlet H_s .

Keywords: Higgs bosons; CP violation; Non Abelian permutational symmetry $S(3)$; 3HDM

1. INTRODUCTION

It is well known that discrete symmetries, such as charge inversion (C), parity (P) and time reversal (T), are useful in understanding elementary particle interactions. These symmetries are preserved in strong and electromagnetic processes [7, 8, 17] but violated in weak interactions as it has been proved since the discovery of CP violation in neutral kaon decays in 1964 [5].

Journal of Nuclear
Physics, Material
Sciences, Radiation and
Applications
Vol-4, No-1,
August 2016
pp. 221–239

Barradas-Guevara, JE
Félix-Beltrán, O
Rodríguez-Jáuregui, E

The Standard Model (SM) of electroweak physics is an $SU(2)_L U(1)_Y$ gauge theory coupled to quarks, leptons and one complex hypercharge-one, $SU(2)_L$ doublet of scalar fields [10, 17]. As it is known in the SM there are three quark families, as first proposed in the theory of CP violation by M. Kobayashi and T. Maskawa [11], based on phases in the coupling constants describing the weak charge-changing transitions of quarks, CP-violating effects arise when there are at least three quark families. For three quark families these couplings are described by the unitary 3×3 Cabibbo-Kobayashi-Maskawa (CKM) matrix [3, 11]. With three quark families, one cannot make a phase redefinition of quarks so that all the couplings are real. To return to the CKM matrix, we have that a unitary 3×3 matrix can be parameterized with three mixing angles θ_{ij} and one CP violating phase δ in the standard parameterization as suggested by M. Kobayashi and T. Maskawa [12]; with $s_{ij} = \sin \theta_{ij}$, $c_{ij} = \cos \theta_{ij}$, and δ is the phase responsible for all CP-violating phenomena in flavor-changing processes in the SM. The angles θ_{ij} can be chosen to lie in the first quadrant, so $s_{ij}, c_{ij} \geq 0$. The unitarity of the CKM matrix, $V^\dagger V = 1$, is the modern way to search for physics beyond the SM. In case of this requirement is violated, it is an indication of the existence of a fourth quark family.

The origin of magnitudes and phases in the CKM matrix is intimately connected with the origin of the quark masses themselves and family replication, in the SM; the physics for the quark masses is related to the Yukawa interactions and the Higgs mechanism. In the basis where the SM Lagrangian is gauge invariant, the charged currents are diagonal and after the electroweak symmetry breaking (EWSB) the quarks, leptons, Higgs fields and gauge bosons acquire mass, but the mass matrices are not diagonal.

Rotating the up and down quark mass matrices to its diagonal form, rotates and mixes the quark fields, and we obtain the corresponding mass eigenvalues; in this basis the charged currents are not diagonal, we obtain the CKM matrix instead.

T.D. Lee noticed that CP violation effects may arise after the EWSB in a model with more than one Higgs. One should keep in mind, in particular, the possibility that there may be a close link between the Higgs sector and CP violation, and one should be open to the possible appearance of $SU(2)$ Higgs doublets of non-standard representations such as singlets and doublets under a flavor symmetry group. Revealing in this way that the full details of the underlying mechanism of electroweak symmetry breaking may be considerably more complex than in the Standard Model.

In the SM due to the form of the scalar potential, one component of the complex scalar field acquires a vacuum expectation value (vev), and the $SU(2)_L \times U(1)_Y$ electroweak symmetry is spontaneously broken down to the $U(1)_{EM}$ gauge symmetry of electromagnetism.

Hermiticity of the Lagrangian requires real SM scalar potential parameter, and the bosonic sector of the electroweak theory is CP-conserving. As we know, the SM with only one Higgs, provides an extremely successful description of the observed electroweak phenomena. Nevertheless, there are a number of motivations to extend the Higgs sector of this model by adding a second and third complex $SU(2)$ doublet of scalar fields.

The theoretical structure of the Standard Model is constrained by the flavor group $S(3)$ ($S(3)$ SM). This model contains three-Higgs doublet fields, leading to numerous relations among Higgs masses and couplings. In particular, as in the case of the SM, in the normal minimum the tree-level $S(3)$ SM Higgs sector is CP-conserving [2, 6].

The $S(3)$ SM Lagrangian contains twelve real scalar fields. After electroweak symmetry breaking, three Goldstone bosons (\mathbf{G}^\pm and \mathbf{G}^0) are removed from the spectrum and provide the longitudinal modes of the massive \mathbf{W}^\pm and \mathbf{Z}^0 . Nine physical Higgs particles remain: two charged Higgs pairs (M_i) and five neutral Higgs. If experimental data reveals the existence of a Higgs sector beyond that of the SM, it will be crucial to test whether the observed scalar spectrum is consistent with an $S(3)$ SM interpretation.

In order to be general within this framework, one should allow for CP-violating arising from $S(3)$ SM when confronting the data. Any observed relations among the $S(3)$ SM parameters would surely contribute to the search for a deeper theoretical understanding of the origin of CP arising from the electroweak symmetry breaking.

In this work we perform a detailed study of the spontaneous CP breaking conditions of $S(3)$ SM. This model has been previously used to calculate the Higgs masses spectrum and mixings as well as trilinear Higgs self-couplings [1], quark and lepton mixing [4, 9], and flavour changing neutral currents (FCNC) [13, 15] with success.

The model has three $S(3)$ flavoured Higgs fields, \mathbf{H}_1 , \mathbf{H}_2 and \mathbf{H}_s , which upon acquiring vev's break the electroweak symmetry. Here, we examine the CP breaking minimization conditions, without explicitly breaking the flavour symmetry, although it might be spontaneously broken. $S(3)$ SM has three different stationary points, which can be classified as Normal, Charge Breaking (CB) and Charge Parity Breaking (CPB) minima, according to the vacuum expectation value of the three Higgs fields [2].

We describe the CPB scenarios of the model and give expressions for the Higgs mass matrix in one CPB scenario. A numerical computation of the trilinear self-couplings $\lambda_{\mathbf{H}_i^0 \mathbf{H}_i^0 \mathbf{H}_i^0}$, $i, \dots, 5$, allows us to find out \mathbf{H}_4^0 , which is the Higgs candidate like to the SM one.

In writing the Higgs potential, we have implicitly chosen a basis in the three-dimensional flavor space of scalar fields. After EWSB, in the CP minimum, the $S(3)$ SM scalar potential depends on real parameters and six Higgs vev's, for a total of sixteen degrees of freedom. The $S(3)$ SM is governed by the choice of the Higgs potential and the Yukawa couplings of the three scalar to the three generations of quarks and leptons. In the gauge basis Φ_a , Φ_b , and Φ_c denote three complex hypercharge-one scalar fields:

$$H_s = \frac{1}{\sqrt{3}}(\Phi_a + \Phi_b + \Phi_c), \begin{pmatrix} H_1 \\ H_2 \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}}(\Phi_a - \Phi_b) \\ \frac{1}{\sqrt{6}}(\Phi_a + \Phi_b - 2\Phi_c) \end{pmatrix}. \quad (1)$$

The quark, lepton and Higgs fields transform in the same way under the flavour symmetry group $S(3)$, all the fields have three species (flavours) and belong to a representation reducible to $1_s \oplus 2$ of $S(3)$. This implies that Q_s and $Q_{1,2}$ for left and right up or down quark fields, and L_s as well as $L_{1,2}$ for left and right charged or neutral lepton fields.

2.2 S(3) Invariant Yukawa Lagrangian

The $S(3)$ SM is governed by the choice of the Higgs potential and the Yukawa couplings of the three scalar for the three generations of quarks and leptons. Let H_1 , H_2 , and H_s denote three complex hypercharge-one, $SU(2)_L$ doublet scalar fields. The challenge is to develop a model that incorporates entirely or partially the CKM mechanism, in this direction, the Yukawa Lagrangian may be written as (Kubo 2003)

$$\begin{aligned} \mathcal{L}_{Y_d} &= -Y_1^d \bar{Q}_I H_s d_{IR} - Y_3^d \bar{Q}_3 H_s d_{3R} \\ &\quad - Y_2^d [\bar{Q}_I k_{IJ} d_{JR} + \bar{Q}_I \eta_{IJ} H_2 d_{JR}] \\ &\quad - Y_4^d \bar{Q} H_s d_{IR} - Y_5^d \bar{Q} H_1 d_{3R} + H.c., \\ \mathcal{L}_{Y_u} &= -Y_1^u \bar{Q}(i\sigma_2) H_s^* u_{IR} - Y_3^u \bar{Q}_3(i\sigma_2) H_s^* u_{3R} \\ &\quad - Y_2^u [\bar{Q}_I k_{IJ}(i\sigma_2) H_1^* u_{JR} + \eta \bar{Q}_I \eta_{IJ}(i\sigma_2) H_s^* u_{JR}] \\ &\quad - Y_4^u \bar{Q}_3(i\sigma_2) H_1^* u_{IR} - Y_5^u \bar{Q}_I(i\sigma_2) H_1^* u_{3R} + H.c. \end{aligned} \quad (2)$$

Singlets carry the index s or 3, and doublets carry indices $I, J = 1, 2$

Higgs Sector with
Spontaneous CP
Violation in $S(3)$
Standard Model

$$k = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \text{ and } \eta = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (3)$$

The $S(3)$ lepton and neutrino invariant Yukawa Lagrangian may also be written by substituting the down quark sector in terms of the charged leptons and up sector by neutral leptons. Furthermore, in this model the mass terms for the Majorana neutrinos may also be introduced,

$$\mathcal{L}_M = -M_1 \nu_{1R}^T C_{\nu_{1R}} - M_3 \nu_{3R}^T C_{\nu_{3R}}, \quad (4)$$

where C is the charged matrix.

2.3 $S(3)$ Invariant Higgs Lagrangian

The Lagrangian \mathcal{L}_H of the $S(3)$ extended Higgs sector incorporates three complex scalar SU(2) doublets fields. Such theory is based on purely flavour symmetry grounds: in view of the family replication of the elementary fermion spectrum one can speculate that this flavour symmetry is the symmetry of the fundamental particles, and an analogous flavour symmetry principle might work for the Higgs sector too:

$$\mathcal{L}_\phi = [D_\mu \phi_S]^2 + [D_\mu \phi_1]^2 + [D_\mu \phi_2]^2 - V(\phi_1, \phi_2, \phi_S), \quad (5)$$

where D_μ is the usual covariant derivative, $D_\mu = \left(\partial_\mu - \frac{i}{2} \mathbf{g}_2 \tau_a W_\mu^a - \frac{i}{2} \mathbf{g}_1 B_\mu \right)$, with \mathbf{g}_1 and \mathbf{g}_2 standing for the U(1) and SU(2) coupling constants.

The only terms that can possibly violate the CP symmetry are the Yukawa couplings and scalar potential. In the SM, the unique source of CP violation comes from the complex phases in the Yukawa couplings that are transferred to the CKM matrix [12] after EWSB. Within such context the possibility of (explicit) CP violation is intimately connected with the presence of a horizontal space: the quarks come in three identical families distinguished only by their masses.

Another source of CP violation could arise in the scalar potential sector [16]. In such case two patterns can be possible, either the CP symmetry is violated explicitly in the theory before EWSB or the CP violation arises spontaneously together with EWSB.

$$H_1 = \begin{pmatrix} \phi_1 + i\phi_4 \\ \phi_7 + i\phi_{10} \end{pmatrix}, H_2 = \begin{pmatrix} \phi_2 + i\phi_5 \\ \phi_8 + i\phi_{11} \end{pmatrix}, H_s = \begin{pmatrix} \phi_3 + i\phi_6 \\ \phi_9 + i\phi_{12} \end{pmatrix}. \quad (6)$$

The subscript s is the flavour index for the Higgs field singlet under $S(3)$. with $i = 1, 2$ are the components of the $S(3)$ doublet field. It is always possible to write the $S(3)$ scalar potential in the following form

$$V(X) = \mathbf{A}^T \mathbf{X} + \frac{1}{2} \mathbf{X}^T \mathbf{B} \mathbf{X}, \quad (7)$$

with the vector $\mathbf{X}^T = (x_1, x_2, x_3, \dots, x_9)$ given by nine real quadratic forms x_i

$$\begin{aligned} x_1 &= H_1^\dagger H_1, & x_4 &= \mathcal{R}e(H_1^\dagger H_2), & x_7 &= \mathcal{I}m(H_1^\dagger H_2), \\ x_2 &= H_2^\dagger H_2, & x_5 &= \mathcal{R}e(H_1^\dagger H_s), & x_8 &= \mathcal{I}m(H_1^\dagger H_s), \\ x_3 &= H_s^\dagger H_s, & x_6 &= \mathcal{R}e(H_2^\dagger H_s), & x_9 &= \mathcal{I}m(H_2^\dagger H_s). \end{aligned} \quad (8)$$

\mathbf{A} is a mass parameter vector,

$$\mathbf{A}^T = (\mu_1^2, \mu_1^2, \mu_0^2, 0, 0, 0, 0, 0, 0) \quad (9)$$

and \mathbf{B} is a 9×9 real parameter symmetric matrix μ_0, μ_1 , and a, b, c, d, e, f, g, h , are real parameters. The $S(3)$ invariant Higgs potential in equation (7) has a CPB minimum at where $\nu_{1,2,3}$ and $\gamma_{1,2,3}$ are real numbers. Here we present the analysis in the case where CPB comes from the Higgs singlet H_s this vacuum solution preserves the $U(1)_{EM}$ symmetry; it corresponds to a local minimum of the potential if its parameters are such that the physical Higgs squared-masses are non-negative.

$$B = \begin{pmatrix} 2(c+g) & 2(c-g) & b & 0 & 0 & 2e & 0 & 0 & 0 \\ 2(c-g) & 2(c+g) & b & 0 & 0 & -2e & 0 & 0 & 0 \\ b & b & 2a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 8g & 4e & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 4e & 2(f+2h) & 0 & 0 & 0 & 0 \\ 2e & -2e & 0 & 0 & 0 & 2(f+2h) & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -8d & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2(f-2h) & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2(f-2h) \end{pmatrix}. \quad \text{Higgs Sector with Spontaneous CP Violation in } S(3) \text{ Standard Model}$$

(10)

$$\phi_7 = v_1, \phi_8 = v_2, \phi_9 = v_3, \phi_{10} = \gamma_1, \phi_{11} = \gamma_2, \phi_{12} = \gamma_3, \text{ and other cases } \phi_i = 0, \quad (11)$$

3. MINIMUM CONDITIONS

In this section, we present the minimum conditions and the parameter space analysis for the considered CPB scenario with $\neq 0$, and $\gamma_1 = \gamma_2 = 0$. Electroweak symmetry breaking arises if the minimum of the scalar potential occurs for nonzero expectation values of the scalar fields.

$$H_i = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_i + i\tilde{a}_i \end{pmatrix} \quad i=1,2,3, \quad (12)$$

where $v_i, \tilde{a}_i \in \mathcal{R}e$. The conditions for extrema of the $S(3)$ scalar potential, considering the CPB conditions of the scalar potential are

$$M_7(\gamma_3) = 2v_1(\gamma_3^2(b+f-2h) + v_3^2(b+f+2h) + 2v_1^2(c+g) + 2v_2^2(c+g) + 6ev_2v_3 + \mu_1^2), \quad (13)$$

$$M_8(\gamma_3) = 2 \left(v_2 \left(\gamma_3^2(b+f-2h) + v_3^2(b+f+2h) + 2v_2^2(c+g) - 3ev_3v_2 + \mu_1^2 \right) + v_1^2 \left(2v_2(c+g) + 3ev_3 \right) \right), \quad (14)$$

Barradas-Guevara, JE
 Félix-Beltrán, O
 Rodríguez-Jáuregui, E

$$M_9(\gamma_3) = 2\left(v_3\left(2a\left(v_3^2 + v_3^2\right) + \mu_0^2\right) + v_1^2\left(v_3(b + f + 2h) + 3ev_2\right) + v_3v_2^2(b + f + 2h) - ev_2^3\right), \quad (15)$$

$$M_{10}(\gamma_3) = 4\gamma_3v_1(ev_2 + 2hv_3), \quad (16)$$

$$M_{11}(\gamma_3) = 2\gamma_3\left(v_2(4hv_3 - ev_2) + ev_1^2\right), \quad (17)$$

$$M_{12}(\gamma_3) = 2\gamma_3\left(2a\left(v_3^2 + v_3^2\right) + \left(v_1^2 + v_2^2\right)(b + f - 2h) + \mu_0^2\right). \quad (18)$$

Here,

$$M_i(\gamma_3) = \frac{\partial V(\phi_1, \phi_2, \dots, \phi_{12})}{\partial \phi_i} \Big|_{\min} = 0$$

with minimum in

$$\min = \{\phi_7 = v_1, \phi_8 = v_2, \phi_0 = v_3, \phi_{12} = \gamma_3, \text{ and other cases } \phi_i = 0\}. \quad (19)$$

From equations (13) and (14), we have

$$v_2M_7 - v_1M_8 = 0.$$

Then,

$$v_1 = \sqrt{3}v_2. \quad (20)$$

Hence, from equation (16), where we take this equal to zero, so that

$$h = -\frac{ev_2}{2v_3}, \quad (21)$$

with these values for v_1 and h from equations (13), (18), we obtain

$$\mu_0^2 = -\frac{av_3(\gamma_3^2 + v_3^2) + 2v_3v_2^2(b+f) + 2ev_2^3}{v_3}, \quad (22)$$

Higgs Sector with
Spontaneous CP
Violation in $S(3)$
Standard Model

$$\mu_1^2 = -\frac{v_3(b+f)(\gamma_3^2 + v_3^2) + 8v_3v_2^2(c+g) + ev_2(\gamma_3^2 + 5v_3^2)}{2v_3}.$$

The minimum constraints are the conditions ensuring that the extremum is a minimum for all directions in ϕ 's space, except for the direction of the Goldstone modes. It is realized when the squared-masses of the nine physical Higgses are all positive. The tree-level amplitudes for the scattering of longitudinal gauge bosons at high energy can be related via the equivalence theorem to the corresponding amplitudes in which the longitudinal gauge bosons are replaced by Goldstone bosons.

We have adopted for convenience $v_i \in \mathcal{R}e$ ($i = 1, 2, 3$). In a standard notation, we can write them as

$$\begin{aligned} v_2 &= v(\sin \omega_3 \sin \omega_{CP})/2, \\ v_3 &= v \cos \omega_3 \sin \omega_{CP}, \\ \gamma_3 &= v \cos \omega_{CP}, \end{aligned} \quad (23)$$

where ω_3, ω_{CP} are two free mixing parameter angles. The scalar potential with spontaneous CP-breaking coming from the Higgs singlet H_s has a minimum in equation (19), which have to satisfy the constraint

$$v = (v_1^2 + v_2^2 + v_3^2 + \gamma_3^2)^{1/2} = 246 \text{ GeV}. \quad (24)$$

We assume Higgs vev's as free parameters subject to the constraint (24), the potential parameters in equation (7), specifically the mass parameters μ_0^2 and μ_1^2 , may be written in terms of the vev's. In the Standard Model (SM), the unique source of CP violation comes from the complex phase in the Cabibbo-Kobayashi-Maskawa (CKM) matrix, which may be transferred from the Yukawa couplings after spontaneous electroweak symmetry breaking (EWSB). For explicit CP violation the Yukawa couplings may all be complex. The possibility of (spontaneous) CP violation is intimately connected with the presence of a horizontal space: the quarks come in three identical families distinguished only by their masses as well as the leptons and Higgs fields as it is assumed in the $S(3)$ SM. In particular for real Yukawa

Barradas-Guevara, JE couplings the corresponding Yukawa Lagrangian is given in [1, 14]. From
Félix-Beltrán, O this, we can express the fermionic mass matrix M_f including spontaneous
Rodríguez-Jáuregui, E CP violation ($\gamma_3 \neq 0$) as

$$M_f = \begin{pmatrix} m_1^{CP} + m_6 & m_2 & m_5 \\ m_2 & m_1^{CP} - m_6 & m_8 \\ m_4 & m_7 & m_3^{CP} \end{pmatrix}, \quad (25)$$

where the CP breaking phase is

$$m_1^{CP} = m_1 - Y_1^f (i\gamma_3), \quad (26)$$

$$m_3^{CP} = m_3 - Y_3^f (i\gamma_3). \quad (27)$$

The remaining m_i ($i=1, 2, \dots, 8$), are the same as in the CP conserving minimum [14]. Thus, the EWSB mechanism provides a source for CP violation in the fermion sector contributing with a CP phase in the quark and lepton mixing matrices. In next section we compute the Higgs mass matrix in the CPB minimum equation (19), the corresponding Higgs mass eigenvalues are given as well.

4. HIGGS MASSES

If we want to know the Higgs mass matrix \mathcal{M}_H^2 , it is necessary to compute the second derivatives of the Higgs potential, equation (7), at the CPB minimum conditions.

$$\left(\mathcal{M}_H^2\right)_{ij} = \frac{1}{2} \frac{\partial^2 V}{\partial \phi_i \partial \phi_j} \Bigg|_{\min} \quad \text{with } i, j = 1, 2, \dots, 12. \quad (28)$$

For three Higgs fields H_1 , H_2 and H_s , equation (6), the corresponding \mathcal{M}_H^2 is a 12×12 real matrix. The Higgs mass matrix is block diagonal with 6×6 symmetric and Hermitian sub-matrices corresponding to the electrically charged $\mathcal{M}_{C,\gamma}^2$ and the neutral $\mathcal{M}_{N,\gamma}^2$ bosons mass matrix respectively. The charged Higgs mass matrix is given by

$$\mathcal{M}_{C,\gamma}^2 = \begin{pmatrix} \mathcal{M}_{C11}^2(\gamma) & \mathcal{M}_{C12}^2(\gamma) \\ -\mathcal{M}_{C12}^2(\gamma) & \mathcal{M}_{C11}^2(\gamma) \end{pmatrix}, \quad (29)$$

Higgs Sector with
Spontaneous CP
Violation in $S(3)$
Standard Model

the mass sub-matrices for charged Higgs in equation (29) are given by

$$\mathcal{M}_{C11}^2(\gamma_3) = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{12} & a_{22} & a_{23} \\ a_{13} & a_{23} & a_{33} \end{pmatrix}, \quad (30)$$

where

$$\begin{aligned} a_{11} &= -2gv_2^2 + \frac{e(3v_3^2 + \gamma_3^2)v_2}{2v_2} + \frac{f(v_3^2 + \gamma_3^2)}{2}, \\ a_{12} &= \sqrt{3}v_2(2gv_2 + ev_3), \\ a_{13} &= \frac{1}{2}\sqrt{3}v_2(ev_2 + fv_3), \\ a_{22} &= -6gv_2^2 + \frac{e(7v_3^2 + \gamma_3^2)v_2}{2v_3} + \frac{f(v_3^2 + \gamma_3^2)}{2}, \\ a_{23} &= \frac{1}{2}v_2(ev_2 + fv_3), \\ a_{33} &= \frac{2v_2^2(2ev_2 + fv_3)}{v_3}. \end{aligned}$$

Then,

$$\mathcal{M}_{C12}^2(\gamma_3) = \begin{pmatrix} 0 & 0 & \frac{1}{2}\sqrt{3}v_2 \left(f + \frac{ev_2}{v_3} \right) \gamma_3 \\ 0 & 0 & \frac{1}{2}v_2 \left(f + \frac{ev_2}{v_3} \right) \gamma_3 \\ -\frac{1}{2}\sqrt{3}v_2 \left(f + \frac{ev_2}{v_3} \right) \gamma_3 & -\frac{1}{2}v_2 \left(f + \frac{ev_2}{v_3} \right) \gamma_3 & 0 \end{pmatrix}. \quad (31)$$

Now, we substituted the equations (30) and (31) in (29) and diagonalize the resulting matrix. The eigenvalues for the charged Higgs fields are degenerated as follows

Barradas-Guevara, JE
Félix-Beltrán, O
Rodríguez-Jáuregui, E

$$\left\{ 0, \frac{(\tilde{a}_3^2 + 4v_2^2 + v_3^2)(ev_2 + fv_3)}{2v_3}, -\frac{ev_2(\tilde{a}_3^2 + 9v_3^2) + fv_3(v_3^2 + \tilde{a}_3^2) + 16gv_3v_2^2}{2v_3} \right\} \quad (32)$$

That is, there are two massless and four non-massless Higgs fields:

$$M_{H_1^\pm} = -\frac{v^2(ev_2 + fv_3)}{2v_3}, \quad (33)$$

$$M_{H_2^\pm} = -\frac{ev_2(\gamma_3^2 + 9v_3^2) + fv_3(v_3^2 + \gamma_3^2) + 16gv_3v_2^2}{2v_3}.$$

The neutral Higgs mass matrix has the form

$$\mathcal{M}_{N,\gamma}^2 = \begin{pmatrix} \mathcal{M}_{N11}^2(\gamma) & \mathcal{M}_{N12}^2(\gamma) \\ \mathcal{M}_{N12}^{2T}(\gamma) & \mathcal{M}_{N22}^2(\gamma) \end{pmatrix}. \quad (34)$$

Here $\mathcal{M}_{N12}^{2T}(\gamma) = (\mathcal{M}_{N12}^2(\gamma))^T$ and the neutral 3×3 Higgs sub-matrices are given by

Hence, we computed the neutral Higgs mass matrix given in equation (34) with equations (35), (36) and (37). Diagonalizing the resulting neutral Higgs mass matrix we obtain: one zero and five non zero eigenvalues: $M_{H_1^0}, \dots, M_{H_5^0}$. In accordance with the SM there is one neutral massless eigenvalue. Then, in the $S(3)$ SM with spontaneous CPB coming from the Higgs singlet H_s there are three Goldstone bosons, one neutral and one charged pair. We also obtained nine massive Higgs fields four electrically charged, with degenerated masses, two by two, and five neutral. The neutral Higgs mass eigenvalues are shown in Figure 1. That is, we have got a physically acceptable final result, because in this spontaneous CPB minimum we have obtained three Goldstone bosons, which can give mass to vector bosons W^\pm and Z^0 , with a massless photon and nine physical Higgs fields, at least one neutral Higgs could have a mass of 125.7 ± 0.4 GeV, and the remaining eight additional Higgs states are candidates for new particles. This scenario provides a strong motivation to extend the analysis to CPB phenomenology arising from spontaneous electroweak symmetry breaking. We denote the masses of these charged Higgs as M_{C_i} and $M_{H_j^0}$ for the neutral Higgs masses, where $i = 1, 2$ and $j = 1, \dots, 5$. In the following

$$\mathcal{M}_{N11}^2 = \begin{pmatrix} 6(c+g)v_2^2 & \sqrt{3}v_2(2(c+g)v_2+3ev_3) & \sqrt{3}v_2(2ev_2+(b+f)v_3) \\ \sqrt{3}v_2(2(c+g)v_2+3ev_3) & 2v_2((c+g)v_2-3ev_3) & v_2(2ev_2+(b+f)v_3) \\ \sqrt{3}v_2(2ev_2+(b+f)v_3) & v_2(2ev_2+(b+f)v_3) & 2av_3^2 - \frac{4ev_2^2}{v_3} \end{pmatrix}, \quad (35)$$

$$\mathcal{M}_{N12}^2(\gamma_3) = \begin{pmatrix} 0 & \sqrt{3}ev_2\gamma_3 & \frac{\sqrt{3}v_2(ev_2+(b+f)v_3)\gamma_3}{v_3} \\ \sqrt{3}ev_2\gamma_3 & -2ev_2\gamma_3 & \frac{v_2(ev_2+(b+f)v_3)\gamma_3}{v_3} \\ -\frac{\sqrt{3}ev_2^2\gamma_3}{v_3} & -\frac{ev_2^2\gamma_3}{v_3} & 2av_3\gamma_3 \end{pmatrix} \quad (36)$$

$$\mathcal{M}_{N22}^2(\gamma_3) = \begin{pmatrix} -\frac{v_2(2(d+g)v_2v_3+e(v_3^2+\gamma_3^2))}{v_3} & \sqrt{3}v_2(2(d+g)v_2+ev_3) & 0 \\ \sqrt{3}v_2(2(d+g)v_2+ev_3) & -\frac{v_2(6(c+g)v_2v_3+e(3v_3^2+\gamma_3^2))}{v_3} & 0 \\ 0 & 0 & 2a\gamma_3^2 \end{pmatrix} \quad (37)$$




Figure 1: The Higgs masses $M_{H_2^\pm}$ and $M_{H_i^0}$ ($i = 1, \dots, 5$) for as a function of the angles where $\pi \leq \omega_{CP} \leq 3\pi/2$ and $-1.9 \leq \omega_{CP} \leq -1.57$. In this region, the Higgs bosons H_j^0 for $j = 1, 4$ are candidates for SM like Higgsboson with mass values at 125.7 ± 10 GeV while $M_{H_{2,3,5}^0}$, will never reach this value. The $M_{H_1^\pm}$ is not shown, and the lower and upper levels of ω_3, ω_{CP} , are set from the masses of the charged Higgs bosons.

section, we analyze the Higgs masses in the spontaneous CPB minimum where $v_1 = \sqrt{3}v_2$.

Higgs Sector with
Spontaneous CP
Violation in $S(3)$
Standard Model

5. $S(3)$ SM PARAMETER SPACE

In this section, we explore γ_3 parameter space regions where the model is consistent. In order to do that, we need to map out regions in the space (v_3, γ_3) where the parameter matrix \mathbf{B} , equation (10), is positive definite, ensuring that we have a CPB minimum point in the $S(3)$ Higgs potential. We assume that the set of quadratic couplings have values where the Higgs masses are $\lesssim 2$ TeV, Table 1. Since M_{H^\pm} only depend of ω_3 , Equation (33), then $-1.9 \leq \omega_3 \leq -1.57$, and from the values for masses de Higgses of Table 1; our numerical analysis is performed using

$$\{a = 10, b = 1, c = 30, d = -6, e = -4.54, f = 6.9, g = 3.5, h < f / 2\};$$

with such values, the matrix \mathbf{B} , equation (10), is positive definite, and the Higgs mass eigenvalues are positive too. Also, these parameter values allowed us to have the mass of the lightest neutral Higgs H_4^0 to be less than 160 GeV.

We can see in Figure 1 the behavior of the masses with respect to the free parameters (ω_3, ω_{CP}) . From equations (24) and (20), we have got equation (23).

The smaller Higgs masses are $M_{H_1^0}$ and $M_{H_4^0}$ while bigger Higgs mass values are allowed for $M_{H_{2,3,5}^0}$ without reaching values greater than 2 TeV. Thus, we evaluate the parameter space and found out values in the range 125.7 ± 0.4 GeV for the Higgs $H_{1,4}^0$, and the remaining one $M_{H_{2,3,5}^0}$ is decoupled from this range, with a mass value running from 300 GeV to 1800 GeV.

The explored ranges for ω_3 and are obtained by considering that $\mu_0^2 < 0$, equation (22), and $\mu_1^2 < 0$. As we remarked before, with the parameter values given in equation (38), the matrix \mathbf{B} , equation (10), is positive definite as well as the CPB Higgs mass eigenvalues. As expected, the explored ω_3 and ω_{CP} parameter range allows heavy neutral Higgses whose masses are greater than 450 GeV, as you can see from Figure 1. Traditionally, in the Higgs potential (7) the quadratic (μ_0^2, μ_1^2) and quartic parameters (a, b, \dots, h) determine the masses of the neutral and bosons charged Higgs. Otherwise, and this is the approach followed here, one can take the free parameters ω_3, ω_{CP} , as input, and determine the parameters of the potential as derived quantities, but, some choices of input will lead to physically acceptable masses, others will not.

Barradas-Guevara, JE **Table1:** Charged and neutral Higgs masses (GeV).
 Félix-Beltrán, O
 Rodríguez-Jáuregui, E

$M_{H_1^\pm}$	$M_{H_2^\pm}$	$M_{H_1^0}$	$M_{H_2^0}$	$M_{H_3^0}$	$M_{H_4^0}$	$M_{H_5^0}$
0-1400	100-900	100-1100	500-955	974-1872	100-634	634-737




Figure 2: The Higgs trilinear self-coupling $\tilde{\lambda}_{H_i^0 H_i^0 H_i^0}$ ($i=1, \dots, 5$), in the space (ω_{CP}, ω_3) , where $\pi \leq \omega_{CP} \leq 3\pi/2$ and $-1.9 \leq \omega_3 \leq -1.57$. Where the intersection is shown in masses $M_{H_i^0}$ with $\tilde{\lambda}_{H_i^0 H_i^0 H_i^0} \sim 1$. The coincidence is given to $M_{H_4^0} \sim 125$ GeV and $\tilde{\lambda}_{H_i^0 H_i^0 H_i^0} \sim 1$, set $\omega_{CP} \sim 3.1$.

The potential (7) is attractive as one $S(3)$ extension of the SM that admits spontaneous CP violation. This is an interesting possibility, since it will become possible to severely constrain or even measure it. From this potential, we can derive as a function of the CP violation parameter \mathcal{V}_3 the Higgses trilinear self-couplings [1], which are shown in Figure 2.

In Figure 1, the neutral Higgs masses with respect to the parameters ω_3 and ω_{CP} are shown, in which CP violation comes from the singlet, equation (6). It can be observed a light Higgs with $M_{H_2^0}, M_{H_4^0}, M_{H_5^0} < 960$ GeV, while the others are < 1800 GeV. Further, one can see that five bosons Higgs find a region in the parameter space such that they reach the values of the masses of 125.0 ± 0.01 GeV, see Figure 2. One can be seen from Figure 2 that each neutral Higgs acquires mass values around 125 GeV in different parameter space regions (ω_3, ω_{CP}) . Then, the computation of the self-couplings allows us to identify a Higgs like the one in the SM. We have to look for parameter space regions (ω_3, ω_{CP}) , such that they simultaneously fit the Higgs mass and trilinear self-coupling for values as in the SM,

$$\tilde{\lambda}_{H_i^0 H_i^0 H_i^0} = \frac{\lambda_{H_i^0 H_i^0 H_i^0}}{\lambda_{h^0 h^0 h^0}^{SM}}, \lambda_{h^0 h^0 h^0}^{SM} = \frac{3M_{h^0}^2}{v}. \quad (39)$$

Figure 2 shows $\tilde{\lambda}_{H_i^0 H_i^0 H_i^0}, i = 1, \dots, 5$, we can identify in this model three possible trilinear self-couplings candidates like to the SM one. Comparing with Figure 1, we found that just H_4^0 is the only SM Higgs candidate in the range 110-140 GeV, with $\tilde{\lambda}_{H_4^0 H_4^0 H_4^0} \sim 1$.

6. CONCLUSIONS

We studied spontaneous CP violation in the most Higgs potential under the non-Abelian flavour symmetry $S(3)$ of the extend SM. Introducing three $SU(2)_L$ Higgs doublet fields in the theory, with twelve real fields. A define structure of the CP violation is obtained which permits the calculation of mass for charge and neutral Higgs boson and we found a parameter space region where the minimum of the potential defines a CPB ground state. A further reduction of free parameters is achieved in the sector scalar, by introducing mixing angles of the critical points of Higgs potential. It can be seen in Figure 2, the Higgs masses in a region of space (ω_3, ω_{CP}) . In this range window, $-1.9 < \omega_3 < -1.57$ and $\pi < \omega_{CP} < 3\pi/2$, $H_{1, \dots, 5}^0$, take values smaller than 2 TeV in agreement observations at this time. The Higgs masses and the trilinear self-coupling are computed in terms only two free parameters. We have shown

Barradas-Guevara, JE
Félix-Beltrán, O
Rodríguez-Jáuregui, E

the Higgs masses and trilinear self-coupling for an allowed parameter set. In this case, the trilinear self-couplings analysis confirms our hypothesis: one can have CP violation resulting from the neutral Higgs sector with a Higgs mass and trilinear self-coupling in accordance with the Standard Model one. Also, we have found a CPN ground state where the CPB comes from the $S(3)$ Higgs singlet. Furthermore, we also computed the Higgs masses and Higgs trilinear self-coupling $\tilde{\lambda}_{H_i^0 H_i^0 H_i^0}, i = 1, \dots, 5$, with ω_3, ω_{CP} . Particularly when

CP violation comes from the $S(3)$ singlet H_s , one observed H_4^0 as a possible candidate like the SM Higgs.

7. ACKNOWLEDGEMENTS

This work has been partially supported *CONACYT-SNI* (México). ERJ acknowledges the financial support received from *FEC* and *PROFOCIE* (México).

REFERENCES

- [1] Barradas-Guevara, E., Felix-Beltran, O., and Rodriguez-Jauregui, E., Trilinear self-couplings in an $S(3)$ flavored Higgs model. *Phys. Rev. D*, **90(9)**, 095001 (2014). <http://dx.doi.org/10.1103/PhysRevD.90.095001>
- [2] Beltran, O. F., Mondragon, M., and Rodriguez-Jauregui, E., Conditions for vacuum stability in an $S(3)$ extension of the standard model. *J. Phys. Conf. Ser.*, **171**, 012028 (2009). <http://dx.doi.org/10.1088/1742-6596/171/1/012028>
- [3] Cabibbo, N., Unitary Symmetry and Leptonic Decays. *Phys. Rev. Lett.*, **10**, 531–533 (1963). <http://dx.doi.org/10.1103/PhysRevLett.10.531>
- [4] Caravaglios, F., Morisi S., hep-ph/0503234 (2005).
- [5] Christenson, J. H., Cronin, J. W., Fitch, V. L., and Turlay, R., Evidence for the 2 pi Decay of the $k(2)0$ Meson. *Phys. Rev. Lett.*, **13**, 138–140 (1964). <http://dx.doi.org/10.1103/PhysRevLett.13.138>
- [6] Emmanuel-Costa, D., Felix-Beltran, O., Mondragon, M., and Rodriguez-Jauregui, E., Stability of the tree-level vacuum in a minimal $S(3)$ extension of the standard model. *AIP Conf. Proc.*, **917**, 390–393 (2007). <http://dx.doi.org/10.1063/1.2751981>
- [7] Glashow, S.L. and Weinberg, S., Natural Conservation Laws for Neutral Currents. *Phys. Rev. D*, **15**, 1958 (1977). <http://dx.doi.org/10.1103/PhysRevD.15.1958>
- [8] Glashow, S., Partial Symmetries of Weak Interactions. *Nucl. Phys.*, **22**, 579–588 (1961). [http://dx.doi.org/10.1016/0029-5582\(61\)90469-2](http://dx.doi.org/10.1016/0029-5582(61)90469-2)
- [9] Gonzalez Canales, et al., Quark sector of $S3$ models: classification and comparison with experimental data. *Phys. Rev. D*, **88**, 096004 (2013). <http://dx.doi.org/10.1103/PhysRevD.88.096004>

-
- [10] Higgs, P. W., Broken Symmetries and the Masses of Gauge Bosons. *Phys. Rev. Lett.*, **13**, 508–509 (1964). <http://dx.doi.org/10.1103/PhysRevLett.13.508>
- [11] Kobayashi, M. and Maskawa, T., CP Violation in the Renormalizable Theory of Weak Interaction. *Prog. Theor. Phys.*, **49**, 652–657 (1973a). <http://dx.doi.org/10.1143/PTP.49.652>
- [12] Kobayashi, M. and Maskawa, T., CP Violation in the Renormalizable Theory of Weak Interaction. *Prog. Theor. Phys.*, **49**, 652–657 (1973b). <http://dx.doi.org/10.1143/PTP.49.652>
- [13] Kubo, J. et al., A minimal $S(3)$ -invariant extension of the standard model. *J. Phys. Conf. Ser.*, **18**, 380–384 (2005). <http://dx.doi.org/10.1088/1742-6596/18/1/013>
- [14] Kubo, J., Mondragon, A., Mondragon, M., and Rodriguez-Jauregui, E., The flavor symmetry. *Prog. Theor. Phys.*, **109**, 795–807 (2003). <http://dx.doi.org/10.1143/PTP.109.795>
- [15] Mondragon, A., Mondragon, M., and Peinado, E., Lepton masses, mixings and FCNC in a minimal $S(3)$ -invariant extension of the Standard Model. *Phys. Rev. D*, **76**, 076003 (2007). <http://dx.doi.org/10.1103/PhysRevD.76.076003>
- [16] Nishi, C., CP violation conditions in N-Higgs-doublet potentials. *Phys. Rev. D*, **74**, 036003 (2006). <http://dx.doi.org/10.1103/PhysRevD.74.036003>
- [17] Weinberg, S., A Model of Leptons. *Phys. Rev. Lett.*, **19**, 1264–1266 (1967). <http://dx.doi.org/10.1103/PhysRevLett.19.1264>
-
- Higgs Sector with Spontaneous CP Violation in $S(3)$ Standard Model